Dr. Scott Trenhaile, MD Michael D. Gilbertson, PAC Tori Homb, APN

SCOTT W. TRENHAILE, M.D. Clinical Assistant Professor/Orthopedic Surgeon

Phone: 815-398-9491 info@scotttrenhailemd.com 324 Roxbury Road | Rockford, Illinois 61107 www.scottrenhailemd.com



Post-Operative Instructions - Partial Menisectomy

Diet

• Following surgery, nausea is very common. Begin with clear liquids and progress to your daily diet as tolerated.

Wound Management

- A bandage is applied to the operative site. Keep this in place for 72 hours after surgery.
- It is normal for there to be drainage and for the dressing to become blood tinged. If this occurs, reinforce with additional dressing.
- Remove the dressing after the 3rd post-operative day. It is still normal for there to some drainage from the incision. Apply steri-strips or band-aids to the incision sites.
- Do not shower until after the dressing is removed. No bathes or hot tubs!
- Do not use any topical agents over the incision sites.
- Sometimes small incisions are made that do not require suture closure. Please do not be alarmed by this.

Activity

- Crutches may be used to assist with walking and you may put as much weight on the operative leg as is comfortable unless otherwise directed by your physician.
- You are encouraged to walk in a controlled environment in order to avoid prolonged sitting,
- Elevate the operative leg when resting with pillows under the foot and ankle to aid in the reduction of swelling.
- You are encouraged to do ankle pumps and quad contractions as soon as you are able.
- No driving until instructed by your physician.
- Sleeping may be done in a reclining chair or bed with the use of pillows to support the operative site as tolerated.
- Generally, if you have a desk job you may return to work on the third post-operative day.
- Return to a physically demanding job will be discussed at your post-operative visit.
- Formal physical therapy will begin after your return visit and it is safe to perform those exercises at home to assist with a return of strength and motion.

Ice/Cryotherapy

• Ice should be used for 30 minute increments 4-5 times a day as tolerated.

If you have any questions during normal clinic hours (8am-5pm), please call Dr. Trenhaile's Nurse – Kari or Clinical Assistant - Alyssa at 815-381-7313.

After hours, please call 815-398-9491.

Page 1 Rev Date: 05/19

Dr. Scott Trenhaile, MD Michael D. Gilbertson, PAC Tori Homb, APN

Phone: 815-398-9491 info@scotttrenhailemd.com 324 Roxbury Road | Rockford, Illinois 61107 www.scottrenhailemd.com





Medication

- Pain Medication: Most patients require the use of narcotic pain medication for a period of time after surgery. You are encouraged to take your prescribed pain medication as indicated on the bottle regardless of pain level for the first 24 hours after surgery.
- Anti-Inflammatory Medication: You may take an over-the-counter anti-inflammatory medication (Ibuprofen or Advil, up to 600mg up to 3 times daily) after surgery in addition to your pain medication, unless you have a history of stomach ulcers. If you are not taking pain medication, you may still take an anti-inflammatory medication for inflammation, pain and swelling. Common side effects of pain medication include: nausea, insomnia, itching, constipation, drowsiness, and hallucinations. Take medications with food to reduce side effects. Do not drive or operative machinery while taking narcotic pain medication.
- Nerve Block: If you received an femoral nerve block, numbness and loss of function may persist for up to 24 hours after surgery.

Post-Operative

• Your post-operative appointment will be 7-10 days after your surgery.

Emergencies

- If at any time you have questions or emergent concerns, contact Dr. Trenhaile, or his Physician Assistant Michael Gilbertson, PA-C at 815-398-9491.
- If you are calling after hours or over a weekend you will be directed to the physician answering service and you may be called by the physician or physician assistant on call.
- Please call if any of the following arise:
 - Fever over 101° (it is normal to have a low grade fever for the first 1-2 days after surgery)
 - Redness
 - Painful swelling
 - Numbness
 - Shortness of breath
 - Excessive bleeding and/or drainage from incision sites
- If you require immediate attention, go to the nearest emergency room